
Regions of star formation as γ -ray sources

Analytic formulae for hadronic production in a single stellar wind

Diego F. Torres¹

Institució de Recerca i Estudis Avançats (ICREA) & Institut de Ciències de l'Espai (IEEC-CSIC), Facultat de Ciències, Universitat Autònoma de Barcelona, Torre C5 Parell, 2a planta, 08193 Barcelona, Spain dtorres@ieec.uab.es

Summary. We present an approximate, analytical formulae for the computation of hadronically-produced gamma-ray luminosity in the innermost regions of stellar winds. We put this into the context of other recent research on regions of star formation as gamma-ray sources.

1 Introduction

It has been proposed that several γ -ray sources are related with early-type stars and their neighborhoods, since collective effects of strong stellar winds as well as of supernova explosions are expected to result in particle acceleration up to energies in excess of a TeV (see, e.g., Montmerle 1979; Cassé & Paul 1980; Bykov 2001, Romero & Torres 2003, Torres et al. 2003). Recently, several TeV unidentified source were detected spatially close to regions of star formation, most especially the first such source, detected in the Cygnus region (Aharonian et al. 2005), where a nearby EGRET source (3EG J2033+4118) has a likely stellar origin (White & Chen 1992; Chen et al. 1996; Romero et al. 1999; Benaglia et al. 2001). The TeV discoveries and the increase in sensitivity at hand in the GeV and TeV regimes has rekindled the interest in regions of star formation as gamma-ray sources. But detecting new populations of gamma-ray sources is not a trivial task, particularly for surveying telescopes. For instance, a large fraction of the detections to be made by the Gamma-ray Large Area Space Telescope (GLAST) will initially be unidentified, and, due to their sheer number, traditional methodological approaches to identify individuals and/or populations of γ -ray sources will encounter procedural limitations. These limitations will hamper our ability to classify source populations lying in the anticipated dataset with the required degree of confidence, particularly those for which no member has yet been convincingly detected in the predecessor experiment EGRET. A new paradigm for achieving the classification of γ -ray source populations based on the implementation

of an a priori protocol to search for theoretically-motivated candidate sources is probably needed in order to protect the discovery potential of the sample (Torres & Reimer 2005). Key to the any new procedure is a statistical assessment by which the discovery of a new population can be claimed under a similar quantitative technique. For the previously cited protocol, this is achieved by introducing that the hypothesis to test in searching for new population of sources is the null hypothesis against a reduced noise level. It is useful to note that the LAT in GLAST will be in a privileged position to actually identify new population of sources. If LAT would have an additional order of magnitude better sensitivity, with no very significant improvement in angular resolution, a situation similar to the GRB case would appear, i.e., a flat distribution of unidentified sources with a few privileged individuals only which are extensively traced in multifrequency studies. Essentially, we would find a γ -ray source coinciding with the position of every member of any population under consideration. And thus, we would lack the capability to achieve discoveries by correlation analysis. This is, perhaps, already indicating that a next generation high energy γ -ray mission after GLAST-LAT should not be exclusively sensitivity-driven if no significant improvement in angular resolution can be achieved.

2 Hadronic processes in single and collective winds

Regions that currently are or have been subject to a strong process of star formation are good candidates to be γ -ray and neutrino emitters and may even be sites where ultra high energy cosmic rays are produced (see Torres & Anchordoqui 2005 for a review). Outside the Galaxy, the more powerful sites of star formation are found within very active galaxies such as starbursts and Luminous or Ultra-Luminous Infrared Galaxies, and they have also been recently analyzed as high energy emitters (e.g. Torres et al. 2003, Torres 2004, Torres et al. 2004ab, Cillis et al. 2005, Domingo-Santamaria & Torres 2005). Here, we will not cover these results concerning collective effects in extragalactic scenarios, referring the reader to the mentioned publications for details. Rather, we will make a short note on the computation of hadronic gamma-rays in single and collective stellar winds.

Typically, a stellar wind consists of an inner region in free expansion, followed by a larger, low dense region of hot shocked wind (bubble), which is finally surrounded by a thin layer of swept-up interstellar medium (ISM) (see, e.g., Lamers & Cassinelli 1999, Ch. 12). We assume that γ -ray production is mainly hadronic and that it occurs at the end of the free expansion phase of the wind evolution, when the swept-up matter is comparable with the mass in the wind driven wave. Its size is then $R_{\text{wind}} = \left(3\dot{M}_\star/4\pi m_p n_0 V_\infty\right)^{1/2}$, where $\dot{M}_\star$ is the star mass loss rate, m_p is the mass of the proton, n_0 is the ISM number density and V_∞ is the wind terminal velocity. For a star of radius

R_* the particle density and particle velocity radial profiles of the wind at this inner zone can be modeled by $n(r) = (1 - R_0/r)^{-\beta} \dot{M}_*/4\pi m_p V_\infty r^2$, and $V(r) = V_\infty (1 - R_0/r)^\beta$, respectively. The parameter β is $\sim 0.8-1$ for massive stars, R_0 is given by $R_0 = R_*(1 - (V_0/V_\infty)^{1/\beta})$, and the wind velocity close to the star, V_0 , is typically $\sim 10^{-2}V_\infty$. As a first exercise, we can compute the resulting γ -ray luminosity of a single star with the target matter being that of the innermost region, and between energies E_1 and E_2 :

$$L_\gamma = \int_{R_*}^{R_{\text{wind}}} \int_{E_1}^{E_2} n(r) q_\gamma(E_\gamma) E_\gamma (4\pi r^2) dr dE_\gamma, \quad (1)$$

with the differential γ -ray emissivity from π^0 -decays, $q_\gamma(E_\gamma)$, approximated by

$$q_\gamma(E_\gamma) = 4\pi\sigma_{pp}(E_p)(2Z_{p \rightarrow \pi^0}^{(\alpha)}/\alpha) J_p(E_\gamma)\eta_A\Theta(E_p - E_p^{\min}) \quad (2)$$

at the energies of interest. The parameter η_A takes into account the contribution from different nuclei in the wind (for a standard composition $\eta_A \sim 1.5$, Dermer 1986). $J_p(E_\gamma)$ is the proton flux distribution evaluated at $E = E_\gamma$ (units of protons per unit time, solid angle, energy-band, and area), which is assumed as a typical power-law $J_p(E_p) = (c/4\pi)K_p E_p^{-\alpha}$, with α being the spectral slope and K_p the spectrum normalization factor. The cross section $\sigma_{pp}(E_p)$ for pp interactions at energy $E_p \approx 10E_\gamma$ can be represented above $E_p \approx 10$ GeV by $\sigma_{pp}(E_p) \approx 30 \times [0.95 + 0.06 \log(E_p/\text{GeV})]$ mb (e.g., Aharonian & Atoyan 1996). $Z_{p \rightarrow \pi^0}^{(\alpha)}$ is the so-called spectrum-weighted moment of the inclusive cross-section. Its value for different spectral indices α is given, for instance, by Drury et al. (1994). Finally $\Theta(E_p - E_p^{\min})$ is a Heaviside function that approximately takes into account the fact that only CRs with energies higher than $E_p^{\min}(r \gg R_*)$ will diffuse into the wind (see below). For normalization purposes, we take the expression of the proton energy density, which is given by

$$\omega_{\text{CR}} = \int N_p(E_p) E_p dE_p = \{134.4; 9.9; 0.8\} K_p \text{ eV cm}^{-3} \equiv \varsigma \omega_{\text{CR}, \odot}, \quad (3)$$

where ς is the enhancement factor of the CR energy density with respect to the local value, $\omega_{\text{CR}, \odot}$, we have considered energies between 1 GeV and 20 TeV, and the choice for the prefactor, $\{134.4; 9.9; 0.8\} \text{ eV cm}^{-3}$, refers to three different slopes, 1.9, 2.0, and 2.1, respectively. If the prefactor is referred as $f(\alpha)$ [with units of $\omega_{\text{CR}, \odot}$, typically given in eV cm^{-3}], the normalization is $K_p(\alpha) \sim \omega_{\text{CR}, \odot} \varsigma / f(\alpha)$, depending on the slope of the spectrum. We assume the Earth-like spectrum to be $J_\oplus(E) = 2.2 E_{\text{GeV}}^{-2.75} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1} \text{ GeV}^{-1}$, so that $\omega_{\text{CR}, \odot} \sim 1.5 \text{ eV cm}^{-3}$. An analytic expression for the γ -ray luminosity of a single star produced in the innermost region of its wind can thus be obtained if the radial dependence of the wind velocity is ignored. This, in fact, does not introduce significant changes on the results because a) the terminal velocity of the wind is reached in only a few star radii, and b) the integration extends

to radii up to $R_{\text{wind}} \gg R_*$, with the largest radii (where most of the mass is located) dominating the sum. The expression for the approximate luminosity then reduces to:

$$L_\gamma \sim \frac{9.6 \times 10^{-38} c \dot{M}_* Z_{p \rightarrow \pi^0}^{(\alpha)} \eta_A K_p(2)}{m_p V_\infty \alpha} (R_{\text{wind}} - R_*) \times \left[0.47 \ln \frac{E_2}{E_1} + \frac{0.06}{2 \log e} \left[\left(\log \left(\frac{E_2}{\text{eV}} \right) \right)^2 - \left(\log \left(\frac{E_1}{\text{eV}} \right) \right)^2 \right] \right] \text{ erg s}^{-1}, \quad (4)$$

for $\alpha=2$, and

$$L_\gamma \sim \frac{9.6 \times 10^{-38} c \dot{M}_* Z_{p \rightarrow \pi^0}^{(\alpha)} \eta_A K_p(\alpha)}{m_p V_\infty \alpha} (R_{\text{wind}} - R_*) \times \left[\frac{0.47}{2 - \alpha} \left(\left(\frac{E_2}{\text{eV}} \right)^{2 - \alpha} - \left(\frac{E_1}{\text{eV}} \right)^{2 - \alpha} \right) - \frac{0.06}{\ln 10 (\alpha - 2)} \left(\left(\frac{E_2}{\text{eV}} \right)^{2 - \alpha} \ln \left(\frac{E_2}{\text{eV}} \right) - \left(\frac{E_1}{\text{eV}} \right)^{2 - \alpha} \ln \left(\frac{E_1}{\text{eV}} \right) \right) - \frac{0.06}{\alpha^2 - 4\alpha + 4} \left(\left(\frac{E_2}{\text{eV}} \right)^{2 - \alpha} - \left(\frac{E_1}{\text{eV}} \right)^{2 - \alpha} \right) \right] \text{ erg s}^{-1}$$

for $\alpha \neq 2$, with the proton mass measured in kg, the terminal velocity and the speed of light measured in cm s^{-1} , the mass loss rate measured in kg s^{-1} , and the wind and stellar radius given in cm. Analogously, the expected γ -ray flux over an energy threshold E_{th} can be evaluated taking into account the distance D from the Earth to the stellar association:

$$F_\gamma(E_\gamma > E_{\text{th}}) = \frac{1}{4\pi D^2} \int_{R_*}^{R_{\text{wind}}} \int_{E_{\text{th}}} n(r) q_\gamma(E_\gamma) 4\pi r^2 dr dE_\gamma. \quad (5)$$

Also here an analytic –although approximate– expression, can be obtained:

$$F_\gamma(E > E_{\text{th}}) \sim \frac{1}{4\pi D^2} \frac{6.0 \times 10^{-26} c \dot{M}_* Z_{p \rightarrow \pi^0}^{(\alpha)} \eta_A K_p}{m_p V_\infty \alpha} (R_{\text{wind}} - R_*) \times \left[\frac{0.47}{\alpha - 1} \left(\frac{E_{\text{th}}}{\text{eV}} \right)^{1 - \alpha} + \frac{0.06}{\ln 10} \left(\frac{E_{\text{th}}}{\text{eV}} \right)^{1 - \alpha} \left(\frac{1}{\alpha^2 - 2\alpha + 1} + \frac{1}{\alpha - 1} \ln \left(\frac{E_{\text{th}}}{\text{eV}} \right) \right) \right] \text{ photons cm}^{-2} \text{ s}^{-1},$$

which is valid for all $\alpha \neq 1$. Here, D must be given in cm, whereas all other quantities must be given in the same units as those used in Eqs. (4) to respect consistency with the numerical factor. This formulae are useful to fix the scale: a typical single star would produce, by interaction between cosmic rays and target matter in the innermost region of its wind, a flux one to three orders of magnitude (depending on the kind of star) below the sensitivity achieved by current telescopes in the TeV regime. Of course, this estimation does not take

into account other processes, particularly those happening in binary systems where two strong stellar winds interact.

In the case of a collection of winds, a hydrodynamical model taking into account the composition of different single stellar winds needs to be adopted. As we see below, differences with the single stellar wind case are notable. One possibility is to consider that there are N stars in close proximity, uniformly distributed within a radius R_c , with a number density $n = 3N/4\pi R_c^3$. Each star has its own mass-loss rate ($\dot{M}_i$) and (terminal) wind velocity (V_i), and average values can be defined as $\dot{M}_w = 1/N \times \sum_i^N \dot{M}_i$, and $V_w = \left(\sum_i^N \dot{M}_i V_i^2 / N \dot{M}_w \right)^{1/2}$. All stellar winds may be assumed to mix with the surrounding ISM and with each other, filling the intra-cluster volume with a hot, shocked, collective stellar wind. A stationary flow in which mass and energy provided by stellar winds escape through the outer boundary of the cluster is established. For an arbitrary distance R from the center of the cluster, mass and energy conservation imply

$$\frac{4\pi}{3} R^3 n \dot{M}_w = 4\pi R^2 \rho V, \quad (6)$$

$$\frac{4\pi}{3} R^3 n \dot{M}_w \left(\frac{1}{2} V_w^2 \right) = 4\pi R^2 \rho V \left(\frac{1}{2} V^2 + h \right), \quad (7)$$

where ρ and V are the mean density and velocity of the cluster wind flow at position R and h is its specific enthalpy (sum of internal energy plus the pressure times the volume). Solving these equations with the corresponding boundary conditions, one obtains the density and velocity of the cluster as a function of distance from its center. This is presented in detail in the paper by Domingo-Santamaría & Torres (2006). Again, it is to be noted that not all cosmic rays will be able to enter the collective wind. The difference between an *inactive target*, as that provided by matter in the ISM, and an *active or expanding target*, as that provided by matter in a single or a collective stellar wind, is indeed given by modulation effects. The cosmic ray penetration into the jet outflow depends on the parameter $\epsilon \sim VR/D$, where V is velocity of wind, and D is the diffusion coefficient. ϵ measures the ratio between the diffusive and the convective timescale of the particles. Taking into account such effects, we have studied collective wind configurations produced by a number of massive stars, and obtained densities and expansion velocities of the stellar wind gas that is target for hadronic interactions in several examples (see details in Domingo-Santamaría & Torres 2006). We have computed secondary particle production, electrons and positrons from charged pion decay, electrons from knock-on interactions, and solve the appropriate diffusion-loss equation with ionization, synchrotron, bremsstrahlung, inverse Compton and expansion losses to obtain expected γ -ray emission from these regions, including in an approximate way the effect of cosmic ray modulation (see Torres & Domingo-Santamaría 2005 for details on the code). Examples where different stellar configurations can produce sources for GLAST and the

MAGIC/HESS/VERITAS telescopes in non-uniform ways, i.e., with or without the corresponding counterparts were found. Cygnus OB 2 and Westerlund 1 maybe two associations where this scenario could be at work

Acknowledgments

DFT has been supported by Ministerio de Educación y Ciencia (Spain) under grant AYA-2006-00530, and the Guggenheim Foundation.

References

1. Aharonian F. A. et al. 2005, A&A 31, 197
2. Aharonian F. A. & Atoyan, A. M. 1996, A&A 309, 91
3. Benaglia P., et al. 2001 A&A 366, 605
4. Bykov, A. M. 2001, Space Sci. Rev., 99, 317
5. Cassé, M. & Paul, J. A. 1980 ApJ, 237, 236
6. Chen, W., White, R. L., & Bertsch, D. 1996, A&AS, 120, 423
7. Cillis A. N., Torres D. F. & Reimer O. 2005, ApJ 621, 139
8. Domingo-Santamaría E. & Torres D. F. 2005, A&A 444, 403
9. Domingo-Santamaría E. & Torres D. F. 2006, A&A 448, 613
10. Lamers H. J. G. L. M., & Cassinelli, J. P. 1999, Introduction to Stellar Winds, Cambridge University Press, Cambridge
11. Montmerle T. 1979, ApJ, 231, 95
12. Romero G. E., & Torres, D. F. 2003, ApJ, 586, L33
13. Romero, G. E., Benaglia, P., & Torres, D. F. 1999, A&A, 348, 868
14. Torres D. F., et al. 2003, Physics Reports, 382, 303
15. Torres D. F. 2004, ApJ 617, 966
16. Torres D. F. & Anchordoqui L. A. 2004, Rept. Prog. Phys. 67, 1663
17. Torres D. F., et al. 2004a, ApJ 607, L99
18. Torres D. F., Domingo-Santamaría E., & Romero G. E. 2004b, ApJ 601, L75
19. Torres D. F. & Reimer O. 2005, ApJ 629, L41
20. Torres D. F. & Domingo-Santamaría E. 2005 Mod. Phys. Lett. A20, 2827
21. White, R. L., & Chen W. 1992, ApJ, 387, L81